

Read the label before opening the container.

For full particulars, see enclosed leaflet.

Citri-Fix

South Africa Reg. No: L9468 Act No. 36 of 1947

A plant growth regulator for prevention of abscission of the calyx on citrus fruit.

ACTIVE INGREDIENT:

2,4-D Sodium salt.....25g/l

Product Information Tel No: 072 678 8226

In case of poisoning: 082 446 8946

Expiry Date:

Batch No:

Date of Manufacture:

UN No. 3082

enviro
bio-chem

Registration holder:

Erintrade CC t/a RT Chemicals
Reg. No: CK 2001/036403/23
44 Kerk Street, Lichtenburg
North West, South Africa 2740
Tel: +27 87 231 7261



HARMFUL
SKADELIK



WARNINGS:

- Handle with care.
- Irritating to eyes, nose, throat and skin.
- Keep out of reach of children, uninformed persons and animals.
- Store in a restricted area away from food and feedstuffs in a cool, dry place.
- Do not handle the treated fruit until the spray deposit has dried unless wearing protective clothing.

Although this remedy has been extensively tested under a large variety of conditions the registration holder does not warrant that it will be efficacious under all conditions because the action and effect thereof may be effected by factors such as abnormal conditions; quality of dilution water, compatibility with other substances not indicated on the label and time and accuracy of the application. The registration holder furthermore does not accept responsibility for damage to crops, vegetation, the environment or harm to man or animal or for lack of performance of the remedy concerned due to failure of the user to follow the label instructions or to the occurrence of conditions which could not have been foreseen in terms of the registration. Consult the supplier in event of any

PRECAUTIONS:

- Do not inhale fumes.
- Wear protective clothing (overall, rubber boots, gloves, eye protection) when handling the concentrate.
- Wash hands with soap and water after use.
- In case of accidental skin contact, wash thoroughly with soap and water.
- Wash contaminated clothing.
- Do not eat, drink or smoke whilst mixing or applying or before washing hands and face and change of clothing.
- Prevent contamination of food, feedstuff, eating utensils and drinking water.
- Prevent contamination onto crops, grazing, rivers, dams or areas not under treatment.
- Clean applicator after use and dispose of wash water where it will not contaminate crops, grazing, dams or rivers.
- Rinse the empty container three times with a volume of water equal to a minimum of 10% of that of the container. Add the rinsings to the contents of the mixing tank before destroying the container in the prescribed manner. Destroy empty container by perforating and flattening and never use for any other purpose.

SYMPTOMS OF HUMAN POISONING:

Coughing, dizziness, temporary loss of muscle co-ordination, fatigue and possible nausea.

FIRST AID TREATMENT:

Treat symptomatically as indicated.

EYES: Flush eyes with clean water for at least 15 minutes. Get medical attention.

SKIN: Wash thoroughly with soap and water.

INHALATION: Remove patient to fresh air.

INGESTION: Induce vomiting.

NOTE TO PHYSICIAN:

There is no antidote. Treat symptomatically.

USE RESTRICTIONS:

Use only as indicated. Refer to RECOMMENDED USAGE RESTRICTIONS FOR PLANT PROTECTION PRODUCTS ON SOUTHERN AFRICAN EXPORT CITRUS of the CRI with regards to MRL's for different markets.

DIRECTIONS FOR USE: Use only as directed.

Mixing Instructions:

Drench and fungicide bath: Pre-mix the required quantity of CITRI-FIX with a small volume of clean water and add to drench or fungicide bath.

Wax: Add required quantity of CITRI-FIX to wax while agitating.

Application:

Apply in a drench, fungicide bath or wax.

APPLICATION TABLE:

CITRUS	DOSAGE	REMARKS
Drench and fungicide bath application.	1ℓ/100ℓ of water	REFER TO MRL'S FOR DIFFERENT MARKETS.
Wax	2ℓ/200ℓ of wax	Wax must be agitated continuously.

NOTE: Do not exceed recommended dosages. Apply within 12 hours after picking for best results.